

Review

The Role of Th17 in Neuroimmune Disorders: A Target for CAM Therapy. Part III**Aristo Vojdani¹, Jama Lambert¹ and Gottfried Kellermann²**¹Immunosciences Lab., Inc., Los Angeles, CA 90035 and ²NeuroScience, Inc., Osceola, WI 54020, USA

Abundant research has mapped the inflammatory pathways leading to autoimmunity and neuroinflammatory disorders. The latest T helper to be identified, Th17, through its proinflammatory cytokine IL-17, plays a pathogenic role in many inflammatory conditions. Today, healthcare providers have a wealth of anti-inflammatory agents from which to choose. On one hand, pharmaceutical companies market brand-name drugs direct to the public and physicians. Medical botanical knowledge, on the other hand, has been passed down from generation to generation. The demands for natural healing therapies have brought corresponding clinical and laboratory research studies to elucidate the medicinal properties of alternative practices. With a variety of options, it can be difficult to pinpoint the proper anti-inflammatory agent for each case presented. In this review, the authors highlight a vast array of anti-inflammatory medicaments ranging from drugs to vitamins and from botanicals to innate molecules. This compilation may serve as a guide for complimentary and alternative healthcare providers who need to target neuroinflammation driven by Th17 and its inflammatory cytokine IL-17. By understanding the mechanisms of anti-inflammatory agents, CAM practitioners can tailor therapeutic interventions to fit the needs of the patient, thereby providing faster relief from inflammatory complaints.

Keywords: helper T cells – herbal medicine – inflammatory response – lipoic acid – probiotics

Introduction

Parts I and II of this series detail the destructive Th17 inflammatory pathway. Here, we present ways to intervene. The absence of Th17 cells can lead to a decrease in proinflammatory cytokines. Thus, to achieve the end goal of a balanced immune response, CAM practitioners may target components along the pathway leading from naïve T-cell differentiation to Th17, thereby lowering the levels of Th17 and IL-17. Treatment may include enhancing immunity by administering an agonist, or depressing Th17 cell function by introducing an antagonist of ROR γ t. The CAM protocol of ROR γ t agonist therapy may be applied to cancer patients when up-regulation of

the immune response is needed. Intervention in autoimmune disorders or neurodegenerative diseases such as irritable bowel disorder, arthritis, diabetes and multiple sclerosis may include the use of antagonists to ROR γ t. Current methods, outlined here, include a choice of medicines ranging from pharmaceuticals to the integrative practice of intravenous therapy, and from natural plant extracts to innate molecules. Whether mainstream or alternative in practice, sometimes the most effective intervention includes a combination of these anti-inflammatory agents that is suitable for CAM.

Targeting Th17, ROR γ t using Medications and Pharmaceuticals

General, or specific, medications have been or are being developed to reduce the symptoms of inflammatory diseases. Neuroprotective medications currently used for

For reprints and all correspondence: Aristo Vojdani, Immunosciences Lab., Inc., 822 S. Robertson Boulevard, Suite 312 Los Angeles, CA 90035, USA. Tel: +1-310-657-1077; Fax: +1-310-657-1053; E-mail: drari@msn.com

blocking microglial activation include minocycline or dextromethorphan (1). The anti-inflammatory mechanisms of minocycline (2,3) and dextromethorphan (4,5) are unknown at this time; however, these products have been shown to be effective against neuroinflammation. An antibiotic that inhibits the response to IL-2 and thereby blocks the activation of both T cells and B cells, rapamycin, now called sirolimus, is used to combat autoimmunity and to prevent acute graft rejection (6). To protect intestinal microflora and prevent further inflammatory damage, the use of antibiotics must be used in conjunction with a probiotic program.

Medications are not always favored; some practitioners prefer to incorporate intravenous therapies for their patients with chronic inflammatory disorders. Insulin-like growth factor-1 (IGF-1) is principally produced by the liver and circulates in the blood. Decreasing with age, a lack of IGF-1 is found in degenerative arthritis, septic shock, cardiovascular disease and inflammation of the bowel (7). Treatment with intravenous immunoglobulin (IVIG), a plasma protein replacement therapy, uses immunoglobulins extracted from multiple healthy blood donors, is in use to decrease the severity of autoimmune disease (8,9). However, since IGF-1 treatment may increase growth hormone and growth factor, this should be the last choice (10).

The use of anti-cytokine treatments that target inflammation in autoimmune disorders is gaining popularity. Recombinant soluble TNF receptor, etanercept (Enbrel[®]), binds to TNF and blocks inflammatory activity, while infliximab (Remicade[®]) and adalimumab (Humira[®]) are monoclonal antibodies specific for TNF, thus also binding to TNF and blocking inflammatory activity (11–14). These anti-cytokine therapies are not specific and their long-term use can result in the development of immune complexes (15). Pharmaceutical reviews by Drugs.com, an independent Internet source for consumers and healthcare providers, offer warnings associated with anti-cytokine medications that include drug interactions, suppressed immunity, exacerbation of heart disease and susceptibility to serious infections. These are just a few examples from a laundry list of potential problems. Full investigation into drug interactions, side effects and possible dire consequences of long-term use needs to be performed before the onset of pharmaceutical, anti-cytokine treatment.

Targeting Th17 through the ROR γ t Pathway

Possible targets for the suppression of Th17 production include four components: ROR γ t, TGF- β , IL-6 and NF- κ B. Inhibition of NF- κ B promotes the production of proinflammatory cytokines and can result in an improved balance between pro- and anti-inflammatory cytokines. Inhibiting IL-6 production, which is required

for the induction of ROR γ t, may result in the inhibition of Th17 and the production of IL-17 cytokines. Enhancing the production of TGF- β , as this cytokine can influence transcription FoxP3, can differentiate the T cell into a regulatory T cell (T_{reg}). Therefore, down-regulating ROR γ t, the transcription factor responsible for T-cell differentiation to Th17, will result in the reduction of the number of Th17 cells and therefore their cytokines. Overall, manipulation of the pathway toward the production of IL-17 can result in the inhibition of an inflammatory cascade.

Targeting ROR γ t

Retinoic acid-related orphan receptor (ROR) γ t is the key that unlocks the differentiation of naïve T-cells to effector Th17 cells. ROR γ t is expressed exclusively by immune system cells, double positive CD4⁺CD8⁺ thymocytes (16). Interestingly, both IL-6 and TGF- β are individually capable of expressing small amounts of ROR γ t; however, neither can singly induce IL-17-secreting Th17. According to Batten *et al.* (17), the over-expression of this transcription factor induces production of Th17 and its effector cytokine IL-17; conversely, a deficiency of ROR γ t nearly halts T-cell differentiation to Th17. Therefore, by eliminating the key that unlocks IL-17-expressing Th17 cells, CAM practitioners may turn around the inflammatory response by targeting ROR γ t cells in their patients presenting with neuroimmune inflammatory disorders.

High levels of ROR γ t are present in inflammatory diseases, autoimmune disorders and food allergies. Treatment of immune-mediated diseases characterized by the presence of inflammatory cytokines, such as inflammatory bowel diseases, multiple sclerosis, arthritis, dry eye autoimmune disease and diabetes, may involve ROR γ t antagonists, while administration of agonists to ROR γ t may be helpful in the treatment of infectious diseases and boosting anti-tumor immunity (16,18). By the enhancement of mucosal immunity or the increase in the number of T cells reactive to a specific antigen, ROR γ t-developed Th17 can protect the host from infection.

The proximity of ROR γ t to Th17 in the development of IL-17 makes it a pivotal step in the immune response. When Th17 protection is necessary, the up-regulation of ROR γ t would be beneficial. On the other hand, if chronic inflammation exists, the production of IL-17 should be minimized, and thus the suppression of ROR γ t is desirable.

Targeting Th17, ROR γ t using Complimentary and Alternative Agents

Due to risks associated with long-term use, undesirable side effects and the ineffectiveness of healing properties

exhibited by many pharmaceutical agents, more and more patients suffering from chronic inflammatory disorders are turning to complementary and alternative medicine therapies. CAM practitioners have a wealth of natural alternatives to reduce inflammation. Previous authors of articles from *Evidence-Based Complementary and Alternative Medicine* and other journals have provided an excellent insight into anti-inflammatories, such as Moutan Cortex (root bark of *Paeonia suffruticosa* Andrews) featured by Chun *et al.* (19) and later by Wu and Gu (20); *Perillae fructus* (perilla seed) by Yim *et al.* (21), atractylenolide I (extracted from largehead *atractylodes rhizome*) by Liu *et al.* (22), and Chen and colleagues' (23) examination of Tibetan medicated-bath therapy. Additional published natural anti-inflammatories that have been shown to inhibit the production of proinflammatory cytokines include *Opuntia humifusa* Raf. (24), total flavonoids of *Fructus Chorspondiatidis* (25), *Artemisia annua* L. (26), *Kummerowia striata* (Thunb.) Schindl (27) and a mini review of plant extracts by Talhouk and colleagues (28) focusing on the anti-inflammatory properties of bioflavonoids (Table 1).

Indeed, the antioxidant activity and inflammatory response of flavonoids, or bioflavonoids, are of interest due to their therapeutic uses as anti-allergic, anti-cancer activity, anti-microbial and anti-inflammatory in the prevention or maintenance of various disorders. Natural antioxidants remove damaging free radicals and inhibit other oxidation reactions. By using different routes during inflammatory responses, flavonoids are able to block a range of known molecules, notably NF- κ B, inducible nitric oxide synthase (iNOS), cooxygenase (COX) and 5 lipoxygenase enzymes, resulting in the

down-regulation of inflammatory cytokines such as IL-1 β , IL-6 and TNF- α .

The list of published studies elucidating CAM therapeutic results against inflammatory responses is enormous. In the interest of space, we will confine our review to well-documented, natural substances that have been demonstrated to modulate Th17 or IL-17 as they are implicated in many neuroimmune, autoimmune and allergic disorders.

Targeting Th17, ROR γ t, using Retinoic Acid, Vitamin D₃, Resveratrol and Probiotics

Retinoic acid, the oxidized form of vitamin A, has long been viewed as important to both the innate and the acquired immune systems. The transcription factor ROR γ t, which if overexpressed, induces T-cell differentiation to Th17, belongs to a retinoid nuclear receptor superfamily (17). Kim (29) provides an excellent review of published research data referencing the regulation of regulatory T cells by retinoids. Recently, researchers have recognized retinoic acid's regulation in immune responses (17,30). It was shown to suppress memory cell IFN- γ production and increase IL-4 secretion (17). Mucida and colleagues (30) completed a study that measured the effect of retinoic acid on Th17 cell development both *in vitro* and *in vivo*. In each case, retinoic acid greatly reduced ROR γ t expression, which resulted in a measurable reduction of Th17 mucosal T cells. Further, all-trans retinoic acid (ATRA) has been reported to inhibit Th17 production and to promote FoxP3 expression, thereby affecting the Th17/T_{reg} polarization (17). Retinoic acid's importance to the immune system cannot overshadow its promotion of glial and

Table 1. Natural alternatives for reducing inflammation in autoimmune conditions

Agent	Reference	Result
Moutan cortex	(19)	Reduced concentration of TNF- α , IL-1 β and IL-6 in LPS-stimulated RAW264.7 cells
Moutan cortex	(20)	Increased production of IL-10 and a decrease in level of proinflammatory cytokines
<i>Perillae Fructus</i>	(21)	Suppressed IL-4 production, inhibiting IgE secretion; reduced IL-5 and IL-13; suppressed excess Th2 activity and rebalanced Th1/Th2
Atractylenolide I	(22)	Up-regulate TNF; down-regulate IL-1
Tibetan medicated-bath	(23)	Decreased IL-6 and TNF- α levels
<i>Opuntia humifusa</i> Raf	(24)	Blocked expression of IL-6 from LPS-stimulated RAW264.7 cells
Total Flavonoids of <i>Fructus Chorspondiatidis</i>	(25)	Inhibited NF- κ B signaling pathway
<i>Artemisia annua</i> L.	(26)	Inhibited IL-17A production; reduced mRNA expression of ROR γ t; reduced IL-6 production
<i>Kummerowia striata</i> (Thunb.) Schindl ethanol extract	(27)	Suppressed NF- κ B activation; up-regulated IL-10 and HO-1
<i>Artemisia alba</i>	(28)	Inhibited IL-1-induced NF- κ B and AP-1 activation
Ginger extract from <i>Alpinia gallanga</i> and <i>Zingiber officinale</i>	(28)	Reduced level of mRNA
<i>Longicera japonica</i>	(28)	Inhibited LPS-induced degradation of I- κ B α and induction of NF- κ B p65, TNF- α and inducible NO synthase (iNOS)
<i>Phlebodium decumanum</i>	(28)	Reduced TNF production; inhibited IL-6 production; increased release of TNF-R2 and IL-1 receptor agonist
<i>Scandix austalis</i> extracts	(28)	Inhibited IL-1-induced NF- κ B and AP-1 activation
<i>Tripterygium wilfordii</i>	(28)	Lowered the inflammatory response
<i>Urtica dioica</i> leaf extracts	(28)	Inhibited NF- κ B

neuronal differentiation. Within the CNS (central nervous system), retinoic acid has been shown to enhance axonal regeneration of differentiated retinal ganglion cells and peripheral sensory neurons (31). Therefore, supplementation with low doses of vitamin A may be useful for treatment of neuroimmune disorders.

1 α ,25-DihydroxyvitaminD₃

The active metabolite of vitamin D₃, 1 α ,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃), which can prevent the maturation of dendritic cells resulting in tolerogenic dendritic cells with increased potential to induce regulatory T cells is a proven potent inhibitor of NF- κ B expression. Taher and colleagues (32), using a mouse model of allergic asthma, demonstrated that the 1,25(OH)₂D₃ augmentation of immunotherapy resulted in increased serum levels of TGF- β , and thus prevented the Th2-driven development of asthma manifestations. In a study comprising patients with multiple sclerosis, researchers saw a rise in TGF- β levels following a 6-month supplementation phase of vitamin D (33). Interestingly, there were no changes in TNF- α , IFN- γ and IL-13 levels. Vitamin D₃ acts by binding in signal transduction pathways such as the regulation of intracellular concentrations, protein kinase C activity, phospholipids metabolism and cyclic nucleotide formation (33). CAM practitioners, by specifically targeting an increase of TGF- β , should consider the effectiveness of vitamin D₃, as it has been shown to reduce autoimmune incidences of cardiovascular disease (34), diabetes (35,36), rheumatoid arthritis (37,38) and lupus (39).

Resveratrol

A phytoalexin produced by several plants when under bacterial or fungal attack; resveratrol has become a popular supplement due to the French paradox. It has been suggested that the resveratrol found in the skin of red grapes explains the low incidence of cardiovascular disease in France where wine is consumed almost daily (40). Indeed, immunostimulant resveratrol has been demonstrated to exhibit antioxidant properties and anti-inflammatory characteristics; it stimulates natural killer cell activity and exercises anti-tumor properties through the suppression of the NF- κ B signaling pathway (41). Gonzales and Orlando (42) studied the effects of resveratrol and curcumin on the NF- κ B-mediated cytokine expression in adipocytes. They concluded that both supplements were able to inhibit TNF- α -activated NF- κ B signaling in adipocytes and thereby reduced TNF- α , IL-1 β and IL-6 cytokine expressions. The authors of this study suggest the potential use of resveratrol and curcumin in the development of therapeutic protocols aimed at alleviating the low-level, chronic inflammation often seen in obese patients with a propensity toward

cardiovascular disorders and insulin resistance associated with type 2 diabetes mellitus. In cases of neuroinflammation, the small molecules of resveratrol can easily pass through the blood–brain barrier where they are shown to be neuron-protective. Resveratrol stimulates the production of nicotinamide adenine dinucleotide (NAD), and therefore, is used to slow the progress of amyotrophic lateral sclerosis (43). The neuroimmune protective characteristics of resveratrol make it an attractive and versatile option for CAM practitioners.

Non-Pathogenic Probiotic Bacteria

The human gut is a home to a vast consortium of symbiotic bacteria. Members of this complex microflora metabolize dietary substances, such as plant polysaccharides, that are otherwise indigestible by their human hosts. Interest in the gut flora has led to numerous investigations to demonstrate that there are beneficial and potentially harmful microorganisms in the intestine, and that the one could be used to influence the activities of the other. These findings led to the ‘probiotic’ concept, originally used to describe microbial feed supplements that stimulate the growth of farm animals. Now, the use of live microbes as dietary supplements has been extended to humans (44). The genera most commonly used in probiotic preparations are *Lactobacillus*, *Bifidobacterium*, *Streptococcus*, *Lactococcus* and some fungal strains.

Some of the health benefits which have been claimed for probiotics include the following: improvement of the normal microflora, prevention of infectious diseases and food allergies, reduction of serum cholesterol, anti-carcinogenic activity, stabilization of the gut mucosal barrier, immune adjuvant properties, alleviation of intestinal bowel disease symptoms and improvement in the digestion of lactose in intolerant hosts. The beneficial effects of probiotic bacteria in relation to the augmentation of the immune system and cytokine production were summarized in a mini review by Galdeano *et al.* (44). Pochard and colleagues (45) experimented with lactic acid bacteria and concluded that its administration enhanced the secretion of IL-12 and IFN- γ , thereby modulating the Th1/Th2 balance in cases of allergy. Menard and colleagues (46) determined that metabolites secreted by lactic acid bacteria exerted an anti-TNF- α effect capable of crossing the intestinal barrier, and showed that LPS binding to Th1 cells and NF- κ B was significantly inhibited. Administration of lactic acid bacteria in prebiotic and probiotic forms can now be tailored to the needs of the patient. The influence on the immune system of individual bacterial strains has been conveniently compiled by Ljungh and Wadström (47). An abbreviated listing is shown in Table 2. Lactic acid used in combination with *Saccharomyces boulardii*, a yeast found on lychee fruit, which has proven to be effective

Table 2. Some individual bacterial strains that influence the immune system (47)

Strain	Effect
<i>L. casei</i> Shirota	Induced IL-12 production
Heat-killed <i>L. casei</i> and <i>L. fermentum</i>	Induced TNF- α production
<i>L. paracasei</i> Ncc2461	Induced CD4 ⁺ T-cells to produce IL-10 and TGF- β
<i>L. rhamnosus</i> GG with LPS	Inhibited TNF- α , but not IL-10
<i>L. reuteri</i> + <i>L. brevis</i>	Induced TNF- α , IL-2, IL-1 β
<i>L. rhamnosus</i> + <i>L. acidophilus</i> + <i>B. lactis</i>	Enhanced immunoreactivity of spleen cells and phagocytes
<i>L. acidophilus</i> + <i>L. delbrueckii</i> ssp <i>bulgaricus</i> + <i>B. bifidum</i>	Produced an anti-inflammatory response
<i>L. casei</i> Shirota	Inhibited translocation of NF- κ B and production of IL-6

L., Lactobacillus; B., Bifidobacterium.

in preventing diarrhea and decreasing toxin A and toxin B, provides protection (48). With this information, CAM practitioners may select lactic acid bacteria with *Saccharomyces boulardii* to enhance TGF- β -bearing regulatory cells and inhibit proinflammatory cytokine production. Therefore, *Lactobacillus*, other probiotic bacteria and their antigens may have a promise as new therapeutic agents for the treatment of many neuro-immune disorders that start in the GI tract and manifest in the nervous system.

Targeting NF- κ B

Part II of this series details the pathophysiology of intestinal epithelial tight junction permeability leading to the disruption of blood–brain barriers by Th17 cells, resulting in neuroinflammation. In this scenario, nuclear transcription factor kappa B (NF- κ B) is a powerful instigator of inflammation. Therefore, targeting NF- κ B may dampen neuroinflammation induced by Th17. Target genes of NF- κ B may be divided into different functional groups to reveal a predominance of those with primarily inflammatory functions. Kempe and colleagues (49) classified NF- κ B target genes and found that genes for monocyte, macrophage, neutrophils and T-cell chemoattractant chemokines and cytokines (RANTES, CCL20, CXCL3, CXCL2, CXCL11, CXCL6, MCP-1, IL-8 and GM-CSF) were up-regulated. Through the mitogen-activated protein kinase (MAPK) pathway, the phosphorylation and degradation of the NF- κ B-specific cytoplasmic inhibitor I κ B α and subsequent activation and nuclear translocation of NF- κ B is set in motion (50). The initiation of NF- κ B activation occurs via cytokine signaling, innate or adaptive immune responses or environmental stressors. NF- κ B's key role is the regulation of immune responses to viral antigens or to bacterial antigens as is implicated in the opening of intestinal tight junctions (51,52). Activation of NF- κ B indicates down-stream interplay between the

hypothalamic–pituitary–adrenal (HPA) axis and sympathetic nerve terminals as the endocrine response to psychological stressors and contributes to changes in the activity of the neuroendocrine axis to cellular behavior (49,50).

Once the intersected activation pathways of NF- κ B are stimulated, an increase in certain proinflammatory cytokine levels is evident. This may involve TNF- α , IL-1 β (51), IL-6 (53) and IL-8 (49,52,53). Infections elicit the inflammatory response of IL-1 β , which affects an increase in the expression of adhesion factors on endothelial cells to allow the transmigration of pathogen-fighting leukocytes to sites of infection. IL-1 β is implicated in rheumatoid arthritis (RA) and enhanced intestinal permeability (51,52). Proinflammatory and anti-inflammatory IL-6 is detected in Castelman's disease, chronic lymphadenitis, Basedow's disease, Sjögren's syndrome and RA (54). Secreted by any cells with toll-like receptors involved in the innate immune response, IL-8 functions to recruit neutrophils to phagocytose the antigen. High levels of IL-8 in pregnant women indicate a higher risk of inducing schizophrenia in offspring (55). Each of these cytokines, IL-1 β , IL-6, IL-8 and TNF- α , is capable of promoting the progression of inflammatory, autoimmune or neuroinflammatory disorders (51–55).

A multitude of complementary, and alternative, medicinal extracts affecting NF- κ B have been studied. Recent publications focusing on antioxidants to inhibit the activation of NF- κ B include aged garlic extract (allicin) (56), beta-carotene (57), curcumin (58) and quercetin (59). Jiang and colleagues (60) concluded that lyceum seed oil is able to inhibit NF- κ B through the modulation of NF- κ B expression; while Rajakangas' group (61) studied nuclear translocation and found white currant to be an effective agent that inhibits NF- κ B. This transcription factor plays key roles in a variety of cellular processes ranging from inflammation to cancer, from differentiation to apoptosis and from immune response to proliferation. Acetyl-boswellic acids were used in a study (62) to show that in LPS-stimulated human peripheral monocytes, these molecules down-regulated TNF- α expression through the inhibition of NF- κ B. These findings suggest acetyl-boswellic acids as a tool for the development of novel therapeutic interventions. Therefore, the modulation of NF- κ B and associated cytokines such as IL-1 β , IL6, IL-8 and TNF- α by using one or a combination of different herbal medicines may suppress Th17 and IL-17 production and inhibit neuroinflammation. CAM practitioners may thus consider NF- κ B modulation for therapeutic intervention or maintenance of neuroimmune inflammatory disorders.

Targeting TGF- β

A pleiotropic cytokine secreted by many cells, TGF-1 β , plays vital neuroimmune roles. It can control cell growth,




Figure 1. Degradation of tryptophan through the Kynurenine pathway by indoleamine 2, 3 dioxygenase and its activation by highly inducible proinflammatory cytokines. Modified from Conklin *et al.* (87).

differentiation, inflammation, cell chemotaxis, apoptosis, hematopoiesis and plays a role in the protection of neurons from cell death induced by glutamate excitotoxic chemical hypoxia, apoptosis and oxidative injury (63). Like its partner in the Th17 differentiation pathway, TGF-1 β possesses bipolar characteristics. It is both a neuroprotective factor and a trigger of neuronal cell death (63). This anti-inflammatory cytokine assists in T-cell differentiation. Under conditions of high concentration of TGF- β , which induces the expression of FoxP3, the naïve T-cell has been shown to differentiate into anti-inflammatory T_{reg}, while in the presence of low levels, it will differentiate to proinflammatory Th17 (64–66). Thus, by increasing the level of TGF- β , CAM practitioners may affect T-cell differentiation by polarizing the Th17/T_{reg} ratio toward anti-inflammatory T_{reg} for cases of inflammation and autoimmunity.

Targeting IL-6

Considered both pro- and anti-inflammatory, IL-6 plays a dual role. When secreted by T-cells and macrophages, IL-6 stimulates immune responses to trauma such as burns and tissue damage, and during bouts of exercise, muscle tissue cells will produce IL-6 (67). Additionally, IL-6 can be produced by endothelial cells, retinal pigment epithelial cells, some tumor cells, astrocytes, fibroblasts, keratinocytes, bone marrow stromal cells and activated T and B cells (68–74). In the presence of IL-6, low concentrations of TGF- β induce the expression of ROR γ t resulting in differentiation to Th17 and thus the secretion of volatile IL-17. Because of its inhibitory effects on TNF- α and IL-1 and activation of its interleukin-1

receptor agonist (IL-1RA) and IL-10, IL-6 can be defined as anti-inflammatory. Due to this duplicity, targeting IL-6 can be tricky. A balance of IL-6 is necessary to keep the level of Th17 cells down, while at the same time inhibiting TNF- α and IL-1. The use of bioflavonoids, previously described and summarized in Table 1, to block NF- κ B resulting in the down-regulation of IL-6 can bring down the cytokines involved in the inflammatory response (56–61).

Targeting Th17, ROR γ t using Other Molecules

Innate agents that merit targeting include somatostatin and kynurenine with norepinephrine (NE). Naturally occurring in the body, somatostatin in gastrointestinal mucosa and kynurenines and NE in the nervous system, together these molecules play regulatory roles in inflammatory responses. The mucosal layer of the gastrointestinal tract is the body's first line of defense against environmental stressors. It is somatostatin's ability to reduce inflammation (75–77), which helps to keep the intestinal barrier functioning in its protective role. For this reason, CAM researchers may target somatostatin and kynurenine using tryptophan and hydroxytryptophan for down-regulation of neuroimmune disorders. Through meticulous studies (78,79), researchers have elucidated the arduous pathways, both protective and toxic, of kynurenines and have found that by inhibiting the kynurenine pathway, brain damage from cerebral inflammation is reduced in animal models (see Fig. 1). Natural molecules may be favored in some therapeutic interventions; however, targeting kynurenines, without upsetting the

multiple neurotransmitter pathways they affect, can be very complicated.

Targeting Somatostatin

As regulatory hormone, somatostatin is expressed in the gastrointestinal tract (where nerve endings deliver somatostatin to the epithelial cells), pancreas and regions of the CNS. Classified as an inhibitory hormone, it has been shown to impede proinflammatory responses (80,81). Of particular interest, somatostatin secreted from non-neuronal cells along the digestive tract plays an important role as a mediator during mucosal inflammatory responses after physiological (induced by TNF- α) and pathophysiological (up-regulation of bacteria) stimulations (76). Chowder and colleagues (76) found that somatostatin effectively inhibited spontaneously secreted IL-8 and IL-1 β , as well as secretion initiated by TNF- α or bacterial invasion stimulation. More recently, Zavros and colleagues (77) concluded that circulating levels of somatostatin increased by infusion of IL-4, while IFN- γ suppressed the D-cell release of this peptide during inflammation; thus, Th1 predominant gastritis may be quelled through the up-regulation of somatostatin. Through reduced inflammation, the immunoregulatory function of somatostatin in the gastrointestinal tract may prevent the possibility of neurodegeneration that can occur after chronic intestinal barrier dysfunction.

Products of activated immune cells may exert their influence over the nervous system either by active transport of circulating cytokines into the CNS or via cytokine receptors located on the vagal nerve (88). A key modulator of the sympathetic nervous system is the neurotransmitter norepinephrine (NE). Various levels of this neurotransmitter play an important role in anti- and proinflammatory biological responses. Increased NE, acting centrally, can contribute to the release of anti-inflammatory cytokines, whereas low concentrations support a proinflammatory biological state (89,90). Therefore, it has become crucial to determine neurotransmitter levels, as sub-optimal quantities favor potentially damaging proinflammatory activities. Considerable evidence has given credence to the important role of NE in manipulating adaptive immunity (91). In a study involving mice, NE was required for the production of a Th1 cell-mediated immune response and perhaps further affected the development of Th1 (92). Additionally, it has been proposed that NE is important in increasing the magnitude of a Th1 immune reaction (93). Altogether, these studies indicate that NE may enhance the eventual production of Th1 cells (88). Many studies have implicated IL-6 in contributing a key dose-dependent role in sympathetic nerve activity. Specifically, the combined concentrations of sub-threshold IL-6 and IL-1 β can attenuate sympathetic nerve activity (88). Under stress-induced situations, where optimal NE levels were present,

NE attenuated the release of TNF- α along with IL-6 (91). Taken together not only can IL-6 and TNF- α augment, attenuate or silence in their effects on rate of release of NE (85), but it is decisively clear that NE itself can significantly influence whether a system is more prone to anti- or proinflammatory responses.

NE is responsible for most of the activity within the sympathetic nervous system. It mediates the inhibition of T- and B-cell activations, inhibits IL-2-generated lymphocyte activation and modulates keratinocyte mitogenesis (91,94,95). Low levels of NE promote a proinflammatory state, while the up-regulation of NE favors anti-inflammatory cytokines (96). Interestingly, all kynurenines and resultant intermediates can be measured in blood and urine (78); therefore, targeting kynurenines in conjunction with NE measurements will provide an invaluable therapeutic advantage for healthcare practitioners to achieve the best patient results. In patients with neuroinflammation and autoimmunity, these intervention methods for targeting Th17 ROR γ t cells ranging from pharmaceutical to plant extracts and other molecules with appropriate references are shown in Fig. 2.

Conclusion

During the past 30 years, discussions about the limitations of Th1/Th2 implications for health and disease have resulted in several major revisions of the hypothesis, and hence, the identification of T_{reg} and Th17 cells. Currently, these subsets of CD4⁺ T-helper cells are classified based upon unique cytokine products, signaling pathways and lineage-specific transcription factors. However, recent data argue for more complexity and flexibility in these subsets than was previously assumed (97). As with any model, flexibility and plasticity of the T-cell subset are the subject of intense investigation and enhancement. As was concluded by Wilson *et al.* (98), 'a future challenge will be to determine with greater clarity how specific combinations of epigenetic modifications are established by networks of lineage-specifying transcription factors, and whether, when and how they can later be removed or selectively modified to achieve or alter T_H-lineage specification'.

Thus, based on the best available evidence, the modification of T_H-lineage requires a full understanding of the neuroimmune inflammatory pathway, which is necessary to combat inflammation, autoimmunity and neurodegeneration. In this regard it is vital to understand the significant roles played in neuroinflammation by Th17 and its effector cytokine, IL-17, in conjunction with IL-23, IL-27 and Th1 autoreactive cells.

Although IL-17 is highly pathogenic, as evidenced by its presence in a variety of inflammatory disorders, its initial intent is not destructive. Th17 plays a protective role in host defense against infection, by inducing




Figure 2. Various CAM intervention methods with appropriate references for targeting TH17 ROR γ t cells in autoimmune disorders or neurodegenerative diseases are shown. Possibilities range from pharmaceuticals to natural plant extracts and innate molecules.

chemokine and G-CSF expression from surrounding cells thereby recruiting neutrophils and macrophages to infected tissues (16,99). Within the intestinal mucosa, naturally occurring Th17 cells control a variety of bacterial and fungal infections at the mucosal surfaces (99–102). It is in the presence of chronic inflammation that Th17 becomes destructive and over produces IL-17 (102,103).

Through careful manipulation of the ROR γ t differentiation of naïve T cells into the Th17 cells cascade using the different CAM modalities described in this manuscript, a balanced immune response can be achieved. Research and technology have defined diverse therapeutic options, which have been mentioned above. With a variety of choices available, therapy can be tailored to fit the individual patient's needs. Whether using pharmaceutical or natural treatments, it is important to establish the patient's level of tissue-specific antibodies, status of immune function, including cytokine production before commencing therapy. Such laboratory assessments should be repeated periodically to monitor the effectiveness of the treatment protocol. Alterations may be required to recover from the inflammation inherent in autoimmune, neuroimmune and neurodegenerative diseases and return to a state of optimal health.

More research and publications are needed before a proper meta-analysis of compiled data can be professionally assessed. However, the small number of studies presented in this manuscript shows encouraging experimental evidence, which may inspire the execution of larger, longitudinal research projects in the near future. This additional information could yield detailed cross-comparisons of integrative and CAM therapies for inflammatory and autoimmune disorders.

Acknowledgments

The authors gratefully acknowledge the valuable insight and suggestions contributed by Dr Edwin Cooper toward reviewing this manuscript, as well as for the illustrations created by Mr Joel Bautista.

References

- Goa H-M, Hong J-S. Why neurodegenerative diseases are progressive: uncontrolled inflammation drives disease progression. *Trends Immunol* 2008;29:357–65.
- Lampl Y, Boaz M, Gilad R, Lorberboym M, Dabby R, Rapoport A, et al. Minocycline treatment in acute stroke: an open-label, evaluator-blinded study. *Neurology* 2007;69:1404–10.

3. Tikka TM, Koistinaho JE. Minocycline provides neuroprotection against N-Methyl-D-aspartate neurotoxicity by inhibiting microglia. *J Immunol* 2001;166:7527–33.
4. Wu DC, Jackson-Lewis V, Vila M, Tieu K, Teismann P, Vadseth C, et al. Blockade of microglial activation is neuroprotective in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine administration. *Ann Neurol* 2002;46:598–605.
5. Zhang W, Wang T, Qin L, Gao H-M, Wilson B, Ali SF, et al. Neuroprotective effect of dextromethorphan in the MPTP Parkinson's disease model: role of NADPH oxidase. *FASEB J* 2004;18:589–91.
6. Kopf H, de la Rosa GM, Howard OM, Chen X. Rapamycin inhibits differentiation of Th17 cells and promotes generation of FoxP3+ T regulatory cells. *Int Immunopharmacol* 2007;7:1819–24.
7. Smith GR, Missailidis S. Cancer, inflammation and the AT1 and AT2 receptors. *J Inflamm* 2004;1:3.
8. Anthony RM, Nimmerjahn F, Ashline DJ, Reinhold VN, Paulson JC, Ravetch JV. Recapitulation of IVIG anti-inflammatory activity with a recombinant IgG Fc. *Science* 2008;320:373–6.
9. Nimmerjahn F, Ravetch JV. The anti-inflammatory activity of IgG: the intravenous IgG paradox. *JEM* 2007;204:11–5.
10. Fuchs S, Feferman T, Meidler R, Margalit R, Sicsic C, Wang N, et al. A disease-specific fraction isolated from IVIG is essential for its immunosuppressive effect of IVIG in experimental autoimmune myasthenia gravis. *J Neuroimmunol* 2008;194:89–96.
11. Feldmann M, Maini RN. TNF defined as a therapeutic target for rheumatoid arthritis and other autoimmune diseases. *Nature Med* 2003;9:1245–50.
12. Tobinick EL, Gross H, Weinberger A, Cohen H. TNF-alpha modulation for treatment of Alzheimer's disease: a 6-month pilot study. *Med Gen Med* 2006;8:25.
13. van Deventer S, Sands BE, Anderson FH, Bernstein CN, Chey WY, Feagan BG, et al. Infliximab maintenance therapy for fistulizing Crohn's disease. *N Engl J Med* 2004;350:876–85.
14. Croom KF, McCormack PL. Adalimumab: in plaque psoriasis. *Am J Clin Dermatol* 2009;10:43–50.
15. Emery P, Reginster J-Y, Appelboom T, Bredveld FC, Edelman E, Kekow JM, et al. WHO Collaborating Centre consensus meeting on anti-cytokine therapy in rheumatoid arthritis. *Rheumatology* 2001;40:699–702.
16. Ivanov II, McKenzie BS, Zhou L, Tadokoro CE, Lepelley A, Lafaille JJ, et al. The orphan nuclear receptor ROR γ t directs the differentiation program of proinflammatory IL-17⁺ T helper cells. *Cell* 2006;126:1121–33.
17. Batten M, Li J, Yi S, Kljavin NM, Danilenko DM, Lucas S, et al. Interleukin 27 limits autoimmune encephalomyelitis by suppressing the development of interleukin 17-producing T cells. *Nat Immunol* 2006;7:929–36.
18. Chauhan SK, El Annan J, Ecoiffier T, Goyal S, Zhang Q, Saban DR, et al. Autoimmunity in dry eye is due to resistance of Th17 to Treg suppression. *J Immunol* 2009;182:1247–52.
19. Chun S-C, Jee SY, Lee SG, Park SJ, Lee JR, Kim SC. Anti-inflammatory activity of the methanol extract of Moutan Cortex in LPS-activated Raw264.7 cell. *eCAM* 2007;4:327–33.
20. Wu M, Gu Z. Screening of bioactive compounds from Moutan Cortex and their anti-inflammatory activities in rat synoviocytes. *eCAM* 2007; doi:10.1093/ecam/nem066.
21. Yim Y-K, Lee H, Hong KE, Kim YI, Ko SK, Kim JE, et al. Anti-inflammatory and immune-regulatory effects of subcutaneous *Perillae fructus* extract injections on OVA-induced asthma in mice. *eCAM* 2007; doi:10.1093/ecam/nem118.
22. Liu Y, Zhongwu Jia, Dong Lei, Wang Rui, Qiu G. A randomized pilot study of atractylenolide I on gastric cancer cachexia patients. *eCAM* 2008;5:337–44.
23. Chen H, Shoumura S, Emura S, Isono H. Tibetan medicated-bath therapy may improve adjuvant arthritis in rat. *eCAM* 2007; doi:10.1093/ecam/nem083.
24. Cho JY, Park S-C, Kim T-W, Kim K-S, Song J-C, Kim S-K, et al. Radical scavenging and anti-inflammatory activity of extracts from *Opuntia humifusa* Raf. *J Pharm Pharmacol* 2006;58:113–9.
25. Ao J, Feng H, Xia F. Transforming growth factor and nuclear factor kappa B mediated prophylactic cardioprotection by Total Flavonoids of *Fructus Chorspondiatis* in myocardial ischemia. *Cardiovasc Drugs Ther* 2007;21:235–41.
26. Wang J-X, Tang W, Zhou R, Wan J, Shi L-P, Zhang Y, et al. The new water-soluble artemisinin derivative SM905 ameliorates collagen-induced arthritis by suppression of inflammatory and Th17 responses. *Brit J Pharmacol* 2008;153:1303–10.
27. Tao JY, Zhao L, Huang ZJ, Zhang XY, Zhang SL, Zhang QG, et al. Anti-inflammatory effects of ethanol extract from *Kummerowia striata* (Thunb.) Schindl on LPS-stimulated RAW 264.7 cell. *Inflammation* 2008;31:154–66.
28. Talhouk RS, Karam C, Fostok S, El-Jouni W, Barbour EK. Anti-inflammatory bioactivities in plant extracts. *J Med Food* 2007;10:1–10.
29. Kim CH. Regulation of FoxP3+ regulatory T cells and Th17 cells by retinoids. *Clin Dev Immunol* 2008; doi:10.1155/2008/416910.
30. Mucida D, Park Y, Kim G, Turovskaya O, Scott I, Kronenberg M, et al. Reciprocal T_H17 and regulatory T cell differentiation mediated by retinoic acid. *Science* 2007;317:256–60.
31. Mey J. Retinoic acid as a regulator of cytokine signaling after nerve injury. *Z Naturforsch* 2001;56c:163–76.
32. Taher YA, van Esch BCAM, Hofman GA, Henricks PAJ, van Oosterhout AJM. 1 α ,25-Dihydroxyvitamin D₃ potentiates the beneficial effects of allergen immunotherapy in a mouse model of allergic asthma: role for IL-10 and TGF- β ¹. *J Immunol* 2008; 180:5211–21.
33. Mahon BD, Gordon SA, Cruz J, Cosman F, Cantorna MT. Cytokine profile in patients with multiple sclerosis following vitamin D supplementation. *J Neuroimmunol* 2003;134:128–32.
34. Wang JT, Pencina MJ, Booth SL, Jacques PF, Ingelsson E, Lanier K, et al. Vitamin D deficiency and risk of cardiovascular disease. *Circulation* 2008;117:503–11.
35. Harris SS. Vitamin D in type 1 diabetes prevention. *J Nutr* 2005;135:323–5.
36. Chiu KC, Chu A, Go VL, Saad MF. Hypovitaminosis D is associated with insulin resistance and beta cell dysfunction. *Am J Clin Nutr* 2004;79:820–5.
37. Merlino LA, Curtis J, Mikuls TR, Cerhan JR, Criswell LA, Saag KG. Vitamin D intake is inversely associated with rheumatoid arthritis: results from the Iowa Women's Health Study. *Arthritis Rheum* 2004;50:72–7.
38. Cutolo M, Otsa K, Uprus M, Paolino S, Serio B. Vitamin D in rheumatoid arthritis. *Autoimmun Rev* 2007;7:59–64.
39. Cutolo M, Otsa K. Review: vitamin D, immunity and lupus. *Lupus* 2008;17:6–10.
40. Baur JA, Sinclair DA. Therapeutic potential of resveratrol: the in vivo evidence. *Nat Rev Drug Discov* 2006;5:493–506.
41. Li T, Fan G-X, Wei W, Li T, Yuan Y-K. Resveratrol induces apoptosis, influences IL-6 and exerts immunomodulatory effect on mouse lymphocytic leukemia both in vitro and in vivo. *Int Immunopharmacol* 2007;7:1221–1231.
42. Gonzales AM, Orlando RA. Curcumin and resveratrol inhibit nuclear factor-kappaB-mediated cytokine expression in adipocytes. *Nutr Metab* 2008;5:17–30.
43. Aebischer P, Kato AC. Playing defense against Lou Gehrig's disease. *Sci Am* 2007;297:86–93.
44. Galdeano CM, de Moreno de Le Blanc A, Vinderola G, Bonet MEB, Perdigon G. Proposed model: mechanisms of immunomodulation induced by probiotic bacteria. *Clin Vaccine Immunol* 2007;14:485–92.
45. Pochard P, Gosset P, Grangette C, Andre C, Tonnel AB, Pestel J, et al. Lactic acid bacteria inhibit T_H2 cytokine production by mononuclear cells from allergic patients. *J Allergy Clin Immunol* 2002;110:617–23.
46. Menard S. Lactic acid bacteria secrete metabolites retaining anti-inflammatory properties after intestinal transport. *Gut* 2004; 53:821–8.
47. Ljungh A, Wadström T. Lactic acid bacteria as probiotics. *Curr Issues Intestl Microbiol* 2006;7:73–90.
48. McFarland LV, Surawicz CM, Greenberg RN, Fekety R, Elmer GW, Moyer KA, et al. A randomized placebo-controlled trial of *Saccharomyces boulardii* in combination with standard antibiotics for *Clostridium difficile* disease. *JAMA* 1994;271:1913–18.
49. Kempe S, Kestler H, Lasar A, Wirth T. NF- κ B controls the global proinflammatory response in endothelial cells: evidence for the regulation of a pro-atherogenic program. *Nucleic Acid Res* 2005;33:5308–16.

50. Bierhaus A, Wolf J, Andrassy M, Rohleder N, Humpert PM, Petrov D, et al. A mechanism converting psychosocial stress into mononuclear cell activation. *PNAS* 2003;100:1920–25.
51. Al-Sadi R, Ye D, Dokladny K, Ma TY. Mechanism of IL-1 β -induced increase in intestinal epithelial tight junction permeability. *J Immunol* 2008;180:5653–61.
52. Brandt S, Kwok T, Hartig R, König W, Backert S. NF- κ B activation and potentiation of proinflammatory responses by the *Helicobacter pylori* CagA protein. *PNAS* 2005;102:9300–5.
53. Sun J, Wang XD, Liu H, Xu JG. Ketamine suppresses intestinal NF-kappaB activation and proinflammatory cytokines in endotoxic rats. *World J Gastroenterol* 2004;10:1028–31.
54. Yoshizaki K, Kuritani T, Kishimoto T. Interleukin-6 in autoimmune disorders. *Semin Immunol* 1992;4:155–66.
55. Brown AS, Hooton J, Schaefer CA, Zhang H, Petkova E, Babulas V, et al. Elevated maternal interleukin-8 levels and risk of schizophrenia in adult offspring. *Am J Psychiatry* 2004;161:889–95.
56. Husan N, Mashiat US, Toossi Z, Khan S, Iobal J, Islam N. Allicin-induced suppression of Mycobacterium tuberculosis 85B mRNA in human monocytes. *Biochem Biophys Res Commun* 2007;335:471–6.
57. Guruvayoorappan C, Kuttan G. Beta-carotene inhibits tumor-specific angiogenesis by altering the cytokine profile and inhibits the nuclear translocation of transcription factors in B16F-10 melanoma cells. *Integr Cancer Ther* 2007;6:258–70.
58. Kasinski AL, Du Y, Thomas SL, Zhao J, Sun SY, Khuri FR, et al. Inhibition of IkappaB kinase-nuclear factor-kappaB signaling pathway by 3,5-bis (2-fluorobenzyldene)piperidin-4-one (EF24), a novel monoketone analog of curcumin. *Mol Pharmacol* 2008;74:654–61.
59. Ruiz PA, Braune A, Hölzlwimmer G, Quintanilla-Fend L, Haller D. Quercetin inhibits TNF-induced NF-kappaB transcription factor recruitment to proinflammatory gene promoters in murine intestinal epithelial cells. *J Nutr* 2007;137:1208–15.
60. Jiang YD, Cao J, Dong QZ, Wang SR. Experimental study of anti-atherosclerosis potency by yeacum seed oil and its possible mechanism. *Zhong Yao Cai* 2007;30:372–7.
61. Rajakangas J, Misikangas M, Pääväranta E, Mutanen M. Chemoprevention by white currant is mediated by the reduction of nuclear beta-catenin and NF-kappaB levels in Min mice adenomas. *Eur J Nutr* 2008;47:115–22.
62. Syrovets T, Buchele B, Krauss C, Laumonier Y, Simmet T. Acetyl-boswellic acids inhibit LPS-mediated TNF- α induction in monocytes by direct interaction with I κ B kinases. *J Immunol* 2005;174:498–506.
63. Qian L, Wei SJ, Zhang D, Hu X, Wilson B, El-Benna J, et al. Potent anti-inflammatory and neuroprotective effects of TGF- β 1 are mediated through the inhibition of ERK and p47^{phox}-S_{er}³⁴⁵ phosphorylation and translocation in microglia. *J Immunol* 2008;181:660–8.
64. Monteleone G, Pallone F, MacDonald TT. Interleukin-21: a critical regulator of the balance between effector and regulatory T-cell responses. *Trends Immunol* 2008;29:290–4.
65. Volpe E, Servant N, Zollinger R, Bogiatzi SI, Hupe P, Barillot E, et al. A critical function for transforming growth factor- β , interleukin 23 and proinflammatory cytokines in driving modulating human T_H-17 responses. *Nat Immunol* 2008;9:650–7.
66. Zhou L, Lopes JE, Chong MMW, Ivanov II, Min R, Victora GD, et al. TGF- β -induced Foxp3 inhibits T_H17 cell differentiation by antagonizing ROR γ t function. *Nature* 2008;453:236–40.
67. Febbraio MA, Pedersen BK. Contraction-induced myokine production and release: is skeletal muscle an endocrine organ? *Exerc Sport Sci Rev* 2005;33:114–9.
68. Gauldie J, Richards C, Harnish D, Lansdorp P, Bauma H. Interferon beta 2/B-cell stimulatory factor type 2 shares identity with monocyte-derived hepatocyte-stimulating factor and regulates the major acute phase protein response in liver cells. *Proc Natl Acad Sci USA* 1987;84:7251–55.
69. Wong G, Clark S. Multiple actions of interleukin 6 within a cytokine network. *Immunol Today* 1988;9:137–9.
70. Wong GG, Witek-Giannotti JS, Temple R, Kriz PA, Ferenz C, Hewick RM, et al. Stimulation of murine hemopoietic colony formation by human IL-6. *J Immunol* 1988;140:3040–44.
71. Walther Z, May LT, Sehgal PB. Transcriptional regulation of the interferon-beta 2/B cell differentiation factor BSF-2/hepatocyte-stimulating factor gene in human fibroblasts by other cytokines. *J Immunol* 1988;140:974–7.
72. Roodman GD, Kurihara N, Ohsaki Y, Kukita A, Hosking D, Demulder A, et al. A potential autocrine/paracrine factor in Paget's disease of bone. *J Clin Invest* 1992;89:46–52.
73. Frei K, Malipiero UV, Leist TP, Zinkernagel RM, Schwab ME, Fontana A. On the central nervous system in viral diseases. *Eur J Immunol* 1989;19:689–94.
74. Taga T, Kishimoto T. Interleukin-6. In: Aggarwal BB, Gutterman JU (eds). *Human Cytokines*. Boston, MA: Blackwell Scientific Publications, 1992, 143.
75. Guillermet-Guibert J, Saint-Laurent N, Davenne L, Rochaix P, Cuvillier O, Culler MD, et al. Novel synergistic mechanism for sst2 somatostatin and TNF- α receptors to induce apoptosis: cross-talk between NF- κ B and JNK pathways. *Cell Death Differ* 2007;14:197–208.
76. Chowhry S, Cahalon L, Lahav M, Schor H, Tal R, Bar-Meir S, et al. Somatostatin through its specific receptor inhibits spontaneous and TNF- α -and bacteria-induced IL-8 and IL-1 β secretion from intestinal epithelial cells. *J Immunol* 2000;165:2955–61.
77. Zavros Y, Kao JY, Merchant JL. Inflammation and cancer III. Somatostatin and the innate immune system. *Am J Physiol Gastrointest Liver Physiol* 2004;286:G698–701.
78. Németh H, Robotka H, Toldi J, Vecsei L. Kynurenines in the central nervous system: recent developments. *CNS Agents Med Chem* 2007;7:45–56.
79. Schwarcz R, Pellicciari R. Manipulation of brain kynurenines: glial targets, neuronal effects, and clinical opportunities. *J Pharmacol Exp Ther* 2002;303:1–10.
80. Hoyer D, Bell GI, Berelowitz M, Epelbaum J, Feniuk W, Humphrey PP, et al. Classification and nomenclature of somatostatin receptors. *Trends Pharmacol Sci* 1995;16:86–8.
81. Patel YC. Somatostatin and its receptor family. *Front Neuroendocrinol* 1999;20:157–98.
82. Pace CS, Murphy M, Conant S, Lacy PE. Somatostatin inhibition of glucose-induced electrical activity in cultured rat islet cells. *Am J Physiol* 1977;233:C164–71.
83. Rutter J, Reick M, Wu LC, McKnight SL. Regulation of Clock and NP AS2 DNA binding by the redox state of NAD cofactors. *Science* 2001;293:510–4.
84. Steinman L. Elaborate interactions between the immune and nervous systems. *Nat Immunol* 2004;5:575–81.
85. Mellor AL, Munn DH. IDO expression by dendritic cells: tolerance and tryptophan catabolism. *Nat Rev Immunol* 2004;4:762–74.
86. Taylor MW, Feng GS. Relationship between interferon gamma, indoleamine 2,3-dioxygenase and tryptophan catabolism. *FASEB J* 1991;5:2516–22.
87. Conklin BR, Hsiao EC, Claeysen S, Dumuis A, Srinivasan S, Forsayeth JR, et al. Engineering GPCR signaling pathways with RASSLs. *Nat Methods* 2008;5:673–8.
88. Sanders VM. Interdisciplinary Research: noradrenergic regulation of adaptive immunity. *Brain Behav Immunol* 2006;20:1–8.
89. Bondareff W, Mountjoy CQ, Roth M, Rossor MN, Iversen LL, Reynolds GP, et al. Neuronal degeneration in locus ceruleus and cortical correlates of Alzheimer disease. *Alzheimer Dis Assoc Disord* 1987;1:256–62.
90. Heneka MT, Galea E, Gavrilyuk V, Dumitrescu-Ozimek L, Daeschner J, O'Banion MK, et al. Noradrenergic depletion potentiates beta-amyloid-induced cortical inflammation: implications for Alzheimer's disease. *J Neurosci* 2002;22:2434–42.
91. Kohm AP, Sanders VM. Norepinephrine and beta 2-adrenergic receptor stimulation regulate CD4+ and B lymphocyte functions in vitro and in vivo. *Pharmacol Rev* 2001;54:487–525.
92. Madden KS, Felten SY, Felten DL, Sundaesan PR, Livnat S. Sympathetic neural modulation of the immune system: I. Depression of T-cell immunity in vivo and in vitro following chemical sympathectomy. *Brain Behav Immunol* 1989;3:72–89.
93. Alaniz RC, Thomas SA, Perez-Melgosa M, Mueller K, Farr AG, Palmiter RD, et al. Dopamine beta-hydroxylase deficiency impairs cellular immunity. *Proc Natl Acad Sci USA* 1999;96:2274–78.

94. Farber K, Pannasch U, Kettenmann H. Dopamine and noradrenaline control distinct functions in rodent microglial cells. *Mol Cell Neurosci* 2005;29:128–38.
95. Li Z, Oben JA, Yang S, Lin H, Stafford EA, Soloski MJ. Norepinephrine regulates hepatic innate immune system in leptin-deficient mice with nonalcoholic steatohepatitis. *Hepatology* 2004;40:434–41.
96. O'Connor RA, Prendergast CT, Sabatos CA, Lau CW, Leech MD, Wraith DC, et al. Cutting edge: Th1 cells facilitate the entry of Th17 cells to the central nervous system during experimental autoimmune encephalomyelitis. *J Immunol* 2008;181:3750–54.
97. Haak S, Croxford AL, Kreymborg K, Heppner FL, Pouly S, Becher B, Waisman A. IL-17A and IL-17F do not contribute vitally to autoimmune neuro-inflammation in mice. *J Clin Invest* 2009;119:61–9.
98. Wilson CB, Rowell E, Sekimata M. Epigenetic control of T-helper-cell differentiation. *Nat Rev Immunol* 2009;9:91–105.
99. Weaver CT, Hatton RD, Mangan PR, Harrington LE. IL-17 family cytokines and the expanding diversity of effector T cell lineages. *Annu Rev Immunol* 2007;25:821–52.
100. Bettelli E, Oukka Mohamed, Kuchroo VK. TH17 cells in the circle of immunity and autoimmunity. *Nature Immunol* 2007;8:345–50.
101. Liang SC, Tan XY, Luxenberg DP, Karim R, Dunussi-Joannopoulos K, Collins M, et al. Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *J Exp Med* 2006;203:2271–79.
102. Dong C. T_H17 cells in development: an updated view of their molecular identity and genetic programming. *Nat Rev Immunol* 2008;8:337–348.
103. Yu JJ, Ruddy MJ, Conti HR, Boonnanantanasarn K, Gaffen SL. The interleukin-17 receptor plays a gender-dependent role in host protection against *Porphyromonas gingivalis*-induced periodontal bone loss. *Infect Immun* 2008;76:4206–4213.

Received November 28, 2008; accepted May 22, 2009